

COPPER COMPLEXES AND THEIR USE

5

BACKGROUND OF THE INVENTION

Field of the Invention: The invention relates to copper complexes of phosphorus compounds, to a process for their preparation and to their use in catalytic coupling reactions.

Brief Description of the Prior Art: Coupling reactions catalyzed by copper complexes of aryl halides or arylsulphonates with heteroatom functionalities, for example thiols, amines and amides, have been
10 described in the literature.

Ligands thereof comprising oxygen or phosphorus compounds are frequently used. For example, Venkataraman et al. (Tetrahedron Letters, **2001, 42**, 4791-4793) disclose the use of preformed complexes of copper
15 dibromide and triphenylphosphine to add aryl halides to secondary aromatic amines.

However, disadvantages of the copper complexes of phosphorus compounds known hitherto are the often low chemoselectivity and the
20 narrow spectrum of reactions in which industrially acceptable conversions and conversion rates can be achieved.

There is, therefore, a need to provide copper complexes of phosphorus compounds which are easy to prepare and afford the desired products in
25 good yields in coupling reactions.

SUMMARY OF THE INVENTION

The present invention therefore provides compounds of the formula (I)

"Express Mail" mailing label number EU942980705US


Date of Deposit January 6, 2004

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to the Assistant Commissioner of Patents and Trademarks, Washington, D.C. 20231

Sara Sue Riley

(Name of person mailing paper or fee)


Signature of person mailing paper or fee)



in which

Het¹, Het² and Het³ are each independently absent, or are oxygen or NR¹
 5 where R¹ is C₁-C₈-alkyl, C₅-C₁₈-aryl or C₆-C₁₉-arylalkyl and

B¹ and B² are each independently C₁-C₈-alkyl, C₅-C₁₈-aryl or C₆-C₁₉-
 arylalkyl, or the B¹ and B² radicals together may be a divalent
 radical having a total of 2 to 40 carbon atoms and

10

B³ is C₁-C₈-alkyl, C₅-C₁₈-aryl, C₆-C₁₉-arylalkyl or a radical having a
 total of 2 to 40 carbon atoms and the valency n,

15

X is halide, (C₁-C₈-haloalkyl)carboxylate, (C₁-C₈-alkyl)carboxylate,
 (C₁-C₈-haloalkyl)sulphonate, (C₅-C₁₈-aryl)sulphonate, cyanide,
 optionally fluorinated acetylacetonate, nitrate, oxinate, phosphate,
 carbonate, hexafluorophosphate, tetraphenylborate, tetrakis-
 (pentafluorophenyl)borate or tetrafluoroborate, and

20

p is 0, 1 or 2 and

n is 1, 2 or 3 and

m is 1, 2, 3, 4, 5 or 6.

In the context of the invention, all of the radical definitions, parameters and illustrations above and listed hereinbelow, in general or within areas of preference, i.e. the particular areas and areas of preference too, may be combined as desired.

5

DETAILED DESCRIPTION OF THE INVENTION

Alkyl, alkoxy, alkylene and alkenylene are each independently a straight-chain, cyclic, branched or unbranched alkyl, alkoxy, alkylene and alkenylene radical respectively, and each of the radicals mentioned may optionally also be substituted by C₁-C₄-Alkoxy. The same applies to the nonaromatic moiety of an aralkyl radical.

C₁-C₄-Alkyl is, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl and tert-butyl, C₁-C₈-alkyl is additionally, for example, n-pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, neopentyl, 1-ethylpropyl, cyclohexyl, cyclopentyl, n-hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl, 1-ethyl-2-methylpropyl, 1-ethyl-2-methylpropyl, n-heptyl and n-octyl, C₁-C₁₂-alkyl is further additionally, for example, adamantyl, the isomeric menthyls, n-nonyl, n-decyl and n-dodecyl, and C₁-C₁₈-alkyl is still further additionally, for example, n-octadecyl.

25

C₁-C₄-Alkoxy is, for example, methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, sec-butoxy and tert-butoxy, C₁-C₈-alkoxy is additionally n-pentoxy, 1-methylbutoxy, 2-methylbutoxy, 3-methylbutoxy, neopentoxy, 1-ethylpropoxy, cyclohexoxy, cyclopentoxy, n-hexoxy and n-octoxy, and

C₁-C₁₂-alkoxy is further additionally, for example, adamantoxy, the isomeric menthoxy radicals, n-decoxy and n-dodecoxy.

5 C₁-C₈-Alkylene is, for example, methylene, 1,1-ethylene, 1,2-ethylene, 1,1-propylene, 1,3-propylene, 1,4-butylene, 1,2-cyclohexoxylene and 1,2-cyclopentylene.

10 **Haloalkyl** is in each case independently a straight-chain, cyclic, branched or unbranched alkyl radical which is singly, multiply, or fully substituted by chlorine or fluorine atoms.

C₁-C₈-haloalkyl is, for example, trifluoromethyl, chlorodifluoromethyl, 2,2,2-trifluoroethyl, pentafluoroethyl, nonafluorobutyl, heptafluoroisopropyl and perfluorooctyl.

15

Aryl is in each case independently a heteroaromatic radical having 5 to 18 framework carbon atoms of which no, one, two or three framework carbon atoms per cycle, but at least one framework carbon atom in the entire molecule, may be substituted by heteroatoms selected from the group of
20 nitrogen, sulphur or oxygen, but is preferably a carbocyclic aromatic radical having 6 to 18 framework carbon atoms.

Examples of carbocyclic aromatic radicals having 6 to 18 framework carbon atoms are phenyl, naphthyl, phenanthrenyl, anthracenyl or
25 fluorenyl, and heteroaromatic radicals having 5 to 14 framework carbon atoms of which no, one, two or three framework carbon atoms per cycle, but at least one framework carbon atom in the entire molecule, may be substituted by heteroatoms selected from the group of nitrogen, sulphur or oxygen are, for example, pyridinyl, oxazolyl, benzofuranyl, dibenzofuranyl
30 or quinolinyl.

The carbocyclic aromatic radical or heteroaromatic radical may also be substituted by up to five identical or different substituents per cycle which are selected from the group of free or protected hydroxyl, cyano, chlorine, fluorine, C₁-C₁₂-alkyl, CO(C₁-C₁₂-alkyl), COO(C₁-C₁₂-alkyl), CO(C₅-C₁₈-aryl), COO(C₅-C₁₈-aryl), CON(C₁-C₁₂-alkyl)₂, C₅-C₁₈-aryl, C₁-C₁₂-haloalkyl, C₁-C₁₂-alkoxy, C₁-C₁₂-haloalkoxy, di(C₁-C₈-alkyl)amino or tri(C₁-C₆-alkyl)siloxy.

Arylalkyl is in each case independently a straight-chain, cyclic or branched or unbranched alkyl radical as defined above which may be singly, multiply or fully substituted by aryl radicals as defined above.

The preferred substitution patterns are defined hereinbelow:

Het¹, Het² and Het³ are preferably each independently oxygen or are absent,

B¹ and B² are preferably each independently secondary C₃-C₈-alkyl or tertiary C₄-C₈-alkyl, C₅-C₁₈-aryl or bis(C₅-C₁₈-aryl), or B¹ and B² together are a divalent radical which is selected from the group of 1,2-phenylene, 1,3-phenylene, 1,2-cyclohexyl, 1,1'-ferrocenyl, 1,2-ferrocenyl, 2,2'-(1,1'-binaphthyl) and 1,1'-biphenyl, and the radicals mentioned may optionally be mono- or polysubstituted by cyano, chlorine, fluorine, C₁-C₁₂-alkyl, C₁-C₁₂-haloalkyl, C₁-C₁₂-alkoxy, C₁-C₁₂-haloalkoxy, di(C₁-C₈-alkyl)amino or tri(C₁-C₈-alkyl)siloxy,

B³ is preferably secondary C₃-C₈-alkyl or tertiary C₄-C₈-alkyl, C₅-C₁₈-aryl, C₆-C₁₉-arylalkyl or a radical having a total of from 2 to 40 carbon atoms and the valency n,

X is preferably chloride, bromide or iodide, trifluoromethanesulphonate, trifluoroacetate, methanesulphonate, benzenesulphonate, cyanide, optionally fluorinated acetylacetonate, hexafluorophosphate or tetrafluoroborate, and particular preference is given to chloride, bromide or iodide.

n is preferably 1 or 2,

m is preferably 1 or 2.

10

It has been found that a ratio of copper to phosphorus atoms in the complex, as illustrated by formula (I), can be attained very easily or automatically especially when the ligand contains at least one poly(C₅-C₁₈)aryl or (C₉-C₁₈)aryl structural element or, as a structural element, at least one (C₅-C₁₈)aryl structural element substituted in the ortho-position by secondary or tertiary alkyl, or, in the case of ligands which contain more than one phosphorus atom, B³ is not 1,1'-bisarylene, 2,2'-bisarylene, 1,2-arylene or 1,2-, 1,3- or 1,4-(C₁-C₈)alkylene.

20 In the present embodiment of the invention, formula (I) includes in particular copper-phosphine complexes, copper-phosphonite complexes and copper-phosphite complexes.

Preferred copper-phosphine complexes are, for example, those which contain the following phosphine ligands:

Bis(2-dicyclohexylphosphino)-2'-(N,N-dimethylamino)biphenyl, 2-(dicyclohexylphosphino)biphenyl, 2-(dicyclohexylphosphino)-2'-methylbiphenyl, 2-(di-tert-butylphosphino)biphenyl or 2-(bisdiphenylphosphino)binaphthyl.

30

Preference is given to 2-(di-tert-butylphosphino)biphenyl or 2-(dicyclohexylphosphino)biphenyl.

Particularly preferred copper-phosphine complexes are $[(\mu\text{-Br})_2\{2\text{-(di-tert-butylphosphino)biphenyl}\}_2\text{Cu}_2]$, $[(\mu\text{-trifluoromethanesulphonato})_2\{2\text{-(di-tert-butylphosphino)biphenyl}\}_2\text{Cu}_2]$, $[(\mu\text{-Br})_2\{2\text{-(dicyclohexylphosphino)biphenyl}\}_2\text{Cu}_2]$ and $[(\mu\text{-trifluoromethanesulphonato})_2\{2\text{-(dicyclohexylphosphino)biphenyl}\}_2\text{Cu}_2]$.

10 Preferred copper-phosphonite complexes are, for example, those which contain the following phosphonite ligands:

1,1'-Biphenyl-2-yl dialkyl phosphonites, for example and with preference 1,1'-biphenyl-2-yl dicyclohexyl phosphonite and 1,1'-biphenyl-2-yl di-tert-butyl phosphonite, 3-[(diisopropylphosphino)oxy]phenyl diisopropyl phosphonite, 3-[(di-tert-butylphosphino)oxy]phenyl di-tert-butyl phosphonite, 3-[(diphenylphosphino)oxy]phenyl diphenyl phosphonite or 3-[(dicyclohexylphosphino)oxy]phenyl dicyclohexyl phosphonite, and even greater preference is given to 3-[(diisopropylphosphino)oxy]phenyl diisopropyl phosphonite.

A particularly preferred copper-phosphonite complex is $[(\mu\text{-Br})_2\{2\text{-(di-tert-butylphosphino)biphenyl}\}_2\text{Cu}_2]$.

25 Preferred copper-phosphite complexes are, for example, those which contain the following phosphite ligands:


1,1'-Binaphthyl-2,2'-diyl isopropyl phosphite, 2,4,8,10-tetra-tert-butyl-6-phenoxy-12H-dibenzo[d,g][1,3,2]dioxaphosphocine, and particular

preference is given to 1,1'-binaphthyl-2,2'-diyl isopropyl phosphite and tris(2,4-di-tert-butylphenyl) phosphite.

A particularly preferred copper-phosphite complex is $[(\mu\text{-Br})_2\{1,1'\text{-binaphthyl-2,2'-diyl isopropyl phosphite}\}_2\text{Cu}_2]$.

The copper phosphorus complexes of the formula (I) according to the invention can be prepared, for example, in a manner known per se by reacting phosphorus compounds of the formula (II)

10



in which n, B¹, B², B³, Het¹, Het² and Het³ each have the definitions and areas of preference specified under the formula (I) with compounds of the formula (III)

15



in which X and p each have the definitions and areas of preference specified in the formula (I).

20

In cases in which p = 0, copper powder, for example, can also be used. Preferred compounds of the formula (III) are:

Copper(I) oxide, copper(II) oxide, copper(I) chloride, copper(I) trifluoromethanesulphonate, copper(I) bromide, copper(I) iodide, copper(II)

25

bromide, copper(II) chloride, copper(II) acetate, copper(II) acetylacetonate or mixtures thereof.

5 The molar ratio of phosphorus atoms in compounds of the formula (II) to copper atoms in compounds of the formula (III) in the preparation of complexes of the formula (I) may generally be 10:1 to 0.5:1, preferably 2:1 to 1:1, more preferably 1.2:1 to 1:1.

10 The compounds of the formula (I) can be prepared separately in an inert organic solvent suitable for this purpose, for example tetrahydrofuran, diethyl ether, toluene, xylene, chloroform, dichloromethane, methanol and/or ethanol.

15 The most favourable amount of solvent to be used can be determined by appropriate preliminary experiments.

The compounds of the formula (I) are prepared from the starting compounds of the formulae (II) and (III) described, for example, by simply combining the two starting compounds in solution at room temperature.

20

It is also possible to prepare the compounds of the formula (I) in situ in a catalysis mixture. For this purpose, the compounds of the formula (II) can also be used in the form of phosphonium salts, for example tetrafluoroborates (see also Netherton, M.R.; Fu, G.C.; Organic Letters
25 (2001), 3(26), 4295-429).

The inventive compounds of the formula (I) are suitable in particular for forming carbon-nitrogen, carbon-oxygen and carbon-sulphur bonds, and also for preparing arylalkines.

30

The invention also encompasses catalysts which comprise compounds of the formula (I).

In addition, the invention also encompasses a process for preparing
5 compounds of the formula (IV)



in which

10 n is 1, 2 or 3 and

Ar is a substituted or unsubstituted aromatic radical and

F is oxygen, sulphur, NR^3 , NR^3CO or ethynediyl, where R^3 is hydrogen,
15 C_1 - C_{12} -alkyl, C_5 - C_{18} -aryl or C_6 - C_{19} -arylalkyl and

R^2 is Ar, C_1 - C_{12} -alkyl, C_1 - C_{12} -haloalkyl, C_2 - C_{12} -alkenyl or C_6 - C_{19} -
arylalkyl,

20 which is characterized in that compounds of the formula (V)



in which

25

Ar is as defined above and

Z is chlorine, bromine, iodine, a diazonium salt or sulphonate

30 are reacted with compounds of the formula (VI)



in which

5 F and R² are each as defined above and

the conversion is effected in the presence of base and compounds of formula (I).

10 The areas of preference for compounds of the formulae (IV) to (VI) are defined hereinbelow:

Ar is preferably carbocyclic aromatic radicals having 6 to 24 framework carbon atoms or heteroaromatic radicals having 5 to 24 framework atoms of which no, one, two or three framework atoms per cycle, but at least one framework atom in the entire molecule, are heteroatoms which are selected from the group of nitrogen, sulphur and oxygen. The carbocyclic aromatic radicals or the heteroaromatic radicals may also be substituted by up to five identical or different substituents per cycle which are selected from the group of hydroxyl, chlorine, fluorine, nitro, cyano, free or protected formyl, C₁-C₁₂-alkyl, C₅-C₁₄-aryl, C₆-C₁₅-arylalkyl, -PO-[(C₁-C₈)-alkyl]₂, -PO-[(C₅-C₁₄)-aryl]₂, -PO-[(C₁-C₈)-alkyl](C₅-C₁₄aryl)], tri(C₁-C₈-alkyl)siloxy or radicals of the formula (VIIa-f)

25



30 in which, each independently,

- A is absent or is a C₁-C₈-alkylene radical and
- B is absent or is oxygen, sulphur or NR⁴,
- 5 where R⁴ is hydrogen, C₁-C₈-alkyl, C₆-C₁₅-arylalkyl or C₅-C₁₄-aryl and
- D is a carbonyl group and
- 10 E is R⁵, OR⁵, NHR⁶ or N(R⁶)₂,
- where R⁵ is C₁-C₈-alkyl, C₆-C₁₅-arylalkyl, C₁-C₈-haloalkyl or C₅-C₁₄-aryl and
- 15 R⁶ is in each case independently C₁-C₈-alkyl, C₆-C₁₅-arylalkyl or C₅-C₁₄-aryl, or N(R⁶)₂ together is a cyclic amino radical and
- W is OH, NH₂ or OM where M may be an alkali metal ion, half an equivalent of an alkaline earth metal ion, an ammonium ion or an
- 20 organic ammonium ion,
- Ar is more preferably phenyl, naphthyl, phenanthrenyl, anthracenyl, biphenyl, binaphthyl, fluorenyl, pyridinyl, oxazolyl, thiophenyl, benzofuranyl, benzothiophenyl, dibenzofuranyl, dibenzothiophenyl,
- 25 furanyl, indolyl, pyridazinyl, pyrazinyl, pyrimidinyl, triazolyl and quinolinyl, and the radicals mentioned may also be substituted by no, one, two or three radicals per cycle, each of which is selected independently from the group of fluorine, nitro, cyano, di(C₁-C₄-alkyl)amino, C₁-C₄-alkyl, C₅-C₁₀-aryl, C₁-C₈-fluoroalkyl, C₁-C₈-

fluoroalkoxy, C₁-C₈-alkoxy, CO(C₁-C₄-alkyl), COO-(C₁-C₄)alkyl, -CON(C₁-C₄-alkyl)₂,

- Ar is even more preferably a phenyl radical which may be further substituted by no, one, two or three radicals, each of which is selected independently from the group of nitro, fluorine, cyano, C₁-C₄-alkyl, C₁-C₄-alkoxy, trifluoromethyl, trifluoromethoxy, CO-(C₁-C₄)-alkyl, COO-(C₁-C₄)-alkyl and -CON(C₁-C₄-alkyl)₂,
- n is preferably 1,
- Z is preferably chlorine, bromine or iodine,
- F is preferably oxygen, sulphur, NR³ or ethindyl, where R³ is hydrogen, C₁-C₄-alkyl or C₅-C₁₈-aryl,
- R² is preferably Ar or C₁-C₁₂-alkyl.

For the process according to the invention, the compounds of the formula (I) are generally used in amounts of 0.02 mol% to 10 mol%, preferably 0.1 mol% to 3 mol%, based on the compounds of the formula (IV) used.

The bases used in the process according to the invention are, for example and with preference alkali metal and/or alkaline earth metal carbonates, hydrogencarbonates, alkoxides, phosphates, fluorides and/or hydroxides, and mention should be made in particular of potassium carbonate and/or sodium carbonate, caesium carbonate, caesium hydrogencarbonate, sodium methoxide, potassium tert-butoxide, potassium amylate, caesium fluoride, potassium phosphate and barium hydroxide. Preference is given

to using potassium carbonate, sodium carbonate, caesium carbonate and/or caesium hydrogencarbonate.

Particular preference is given to using potassium carbonate.

5

Per mole of Hal in compounds of the formula (IV) to be exchanged, for example, 0.05 to 10 mol of base can be used, preferably 0.3 to 2 mol.

It is advantageous for the process according to the invention when the
10 bases used are pretreated by grinding and/or drying.

After the grinding, the specific surface areas of the bases are preferably from approx. 0.1 to 10 m²/g, more preferably from 0.2 to 1 m²/g (BET).

15 As a consequence of the marked hygroscopic properties of the bases used in the process according to the invention, the phosphates and carbonates in particular tend to absorb, to a greater or lesser extent, atmospheric constituents such as water and carbon dioxide. From an absorption of approx. 30 per cent by weight of atmospheric constituents, a
20 distinct influence on the conversions to be achieved can be detected. It is therefore frequently appropriate to dry the bases in addition to the grinding.

Depending on the nature of the bases used, the bases are dried, for
25 example, by heating them to temperatures of approx. 50 to 200°C, preferably 100 to 160°C, under a reduced pressure of approx. 0.01 to 100 bar for several hours.

The molar ratio of compounds of the formula (VI) to compounds of the formula (IV) may be, for example, 0.8 to 10, preferably 1 to 6 and more preferably 1 to 4.

- 5 The process according to the invention can be carried out, for example, at temperatures of 20 to 250°C, preferably 100 to 150°C. The optimum reaction temperatures depend in particular on the type of the starting product, of the catalyst and of the bases used, and can be determined by simple preliminary experiments.

10

- The process according to the invention can be carried out either in the presence or in the absence of a suitable solvent. Useful solvents are, for example, aliphatic, alicyclic or aromatic hydrocarbons, for example benzene, toluene, xylene, petroleum ether, hexane, cyclohexane; 15 ethers such as diethyl ether, diisopropyl ether, dioxane, tetrahydrofuran or ethylene glycol dimethyl ether or ethylene glycol diethyl ether; amides, for example N,N-dimethylformamide, N,N-dimethylacetamide, N-methylformanilide, N-methylpyrrolidone or hexamethylphosphoramide; esters such as methyl acetate or ethyl acetate, or mixtures of such solvents.

20

In some cases, an excess of compounds of the formula (VI) can also serve as the reaction medium.

- 25 An azeotroping agent which continuously removes water formed in the course of the reaction azeotropically in the course of the distillation can optionally be added to the process according to the invention.

The process according to the invention can be carried out by customary methods in continuous or batchwise mode.

30

The advantage of the present invention is in particular the easy preparation of the compounds of the formula (I) and the high efficiency with which the inventive copper complexes can be used to prepare compounds of the formula (VI).

5

The invention is further described by the following illustrative but non-limiting examples.

EXAMPLES

10

Preparation of compounds of the formula (I)

Example 1

15 Preparation of bis(2-(di-tert-butylphosphino)biphenylcopper(I) bromide)

50 ml of degassed, anhydrous methanol were heated to reflux temperature, and 2.36 g (7.9 mmol) of 2-(di-tert-butylphosphino)biphenyl
20 were added slowly to the methanol until the phosphine compound was completely dissolved. Subsequently, 0.59 g (2.6 mmol) of copper(II) bromide was added to the solution in portions. After the copper bromide had been added, the solution was heated to reflux temperature for a further 15 min, and then the solution was cooled. After the solution had
25 been cooled, a solid precipitated out and was filtered off, and was washed with a little ethanol and diethyl ether and subsequently dried. 0.93 g (1.1 mmol) of the abovementioned compound was obtained. The yield was 80% of theory.

Exempl 2**Preparation of bis(N-[2'-(dicyclohexylphosphino)-1,1'-biphenyl-2-yl]-N,N-dimethylaminecopper(I) bromide) dimer**

5

50 ml of degassed, anhydrous methanol were heated to reflux temperature, and 2.00 g (5.1 mmol) of N-[2'-(dicyclohexylphosphino)-1,1'-biphenyl-2-yl]-N,N-dimethylamine were added slowly to the methanol until the phosphine compound was completely dissolved. Subsequently, 0.8 g (3.7 mmol) of copper(II) bromide was added to the solution in portions. After the copper bromide had been added, the solution was heated to reflux temperature for a further 15 min, and then the solution was cooled. After the solution had been cooled, a solid precipitated out and was filtered off, and was washed with a little ethanol and diethyl ether and subsequently dried. 1.5 g (1.4 mmol, M = 1073.8 g/mol) of the abovementioned compound were obtained. The structure was checked by an FD-MS analysis (m/e = 1074). The yield was 73% of theory.

10

15

Example 3

20

Preparation of (3-[(diphenylphosphino)oxy]phenyl diphenyl phosphonite)copper(I) chloride

In a round-bottomed flask, 5 g (10.4 mmol) of 3-[(diphenylphosphino)oxy]phenyl diphenyl phosphonite are dissolved in degassed, anhydrous dichloromethane and heated to 40°C. 0.35 g of copper(I) chloride (0.35 mmol) is added. After 30 minutes of stirring, the solvent is removed under reduced pressure. The abovementioned compound was obtained.

25

30

Example 4**Preparation of bis-[(1,1'-biphenyl-2-yl dicyclohexyl phosphonite)copper(I) chloride]**

5

In a round-bottomed flask, 5 g (13.6 mmol) of 1,1'-biphenyl-2-yl dicyclohexyl phosphonite are dissolved in anhydrous, degassed chloroform. In an argon countercurrent, 0.45 g (4.5 mmol) of copper(I) chloride is added and the mixture is stirred at room temperature for 6
10 hours. The solvent is removed under reduced pressure and the residue taken up in anhydrous ether. The mixture is cooled to -78°C and the product precipitates out.

Example 5

15

Preparation of bis[4-isopropoxydinaphtho[2,1-d:1',2'-f][1,3,2]dioxaphosphepine]copper(I) chloride

In a round-bottomed flask, 5 g (13.4 mmol) of 4-isopropoxydinaphtho[2,1-d:1',2'-f][1,3,2]dioxaphosphepine (rac-binaphthyl isopropyl phosphite) are
20 dissolved in dichloromethane and 0.44 g (0.45 mmol) of copper(I) chloride is added in an argon countercurrent. After 30 minutes of stirring, the solvent is removed under reduced pressure.

Examples 6 to 24

25

Use of the catalysts of examples 1 and 2 in coupling reactions

Example 6**Coupling of p-bromoacetophenone with n-octanethiol (catalyst from example 1)**

5

1.35 g (6.7 mmol) of p-bromoacetophenone, 1.0 g (6.7 mmol) of n-octanethiol, 1.8 g (13.5 mmol) of potassium carbonate and 300 mg (0.7 mmol) of the catalyst from example 1 are stirred in 50 ml of dioxane under an argon atmosphere at 110°C for 12 h. The reaction solution is

10 subsequently admixed with 40 ml of aqueous ammonia and extracted with ethyl acetate, and the combined organic extracts are dried under reduced pressure. After workup by column chromatography (hexane), 1.24 g (70%) of product are obtained.

GC-MS/EI: 264 (M)

15

Example 7**Coupling of p-bromoacetophenone with n-octanethiol (catalyst from example 2)**

20

1.35 g (6.7 mmol) of p-bromoacetophenone, 1.0 g (6.7 mmol) of n-octanethiol, 1.8 g (13.5 mmol) of potassium carbonate and 360 mg (0.7 mmol) of the catalyst from example 2 are stirred in 50 ml of dioxane under an argon atmosphere at 110°C for 12 h. The reaction solution is

25 subsequently admixed with 40 ml of aqueous ammonia and extracted with ethyl acetate, and the combined organic extracts are dried under reduced pressure. After workup by column chromatography (hexane), 733 mg (40%) of product are obtained.

Exempl 8**Coupling of p-bromoacetophenone with thiophenol (catalyst from example 1)**

5 1.35 g (6.7 mmol) of p-bromoacetophenone, 0.75 g (6.7 mmol) of thiophenol, 1.8 g (13.5 mmol) of potassium carbonate and 300 mg (0.7 mmol) of the catalyst from example 1 are stirred in 50 ml of dioxane under an argon atmosphere at 110°C for 12 h. The reaction solution is
10 subsequently admixed with 20 ml of aqueous ammonia and extracted with ethyl acetate, and the combined organic extracts are dried under reduced pressure. After workup by column chromatography (hexane), 970 mg (65%) of product are obtained.
GC-MS/EI: 228 (M)

15

Example 9**Coupling of p-bromoacetophenone with thiophenol (catalyst from example 2)**

20 1.35 g (6.7 mmol) of p-bromoacetophenone, 0.75 g (6.7 mmol) of thiophenol, 1.8 g (13.5 mmol) of potassium carbonate and 360 mg (0.7 mmol) of the catalyst from example 2 are stirred in 50 ml of dioxane under an argon atmosphere at 110°C for 12 h. The GC analysis of the
25 crude product indicates product formation with 75% conversion.

Exempl 10**Coupling of p-bromoacetophenone with phenol (catalyst from example 1)**

- 5 1.35 g (6.7 mmol) of p-bromoacetophenone, 630 mg (6.7 mmol) of phenol,
1.8 g (13.5 mmol) of potassium carbonate and 300 mg (0.7 mmol) of the
catalyst from example 1 are stirred in 50 ml of dioxane under an argon
atmosphere at 110°C for 12 h. The reaction solution is subsequently
admixed with 40 ml of aqueous ammonia and extracted with ethyl acetate,
10 and the combined organic extracts are dried under reduced pressure.
After workup by column chromatography (hexane), 280 mg (20%) of
product are obtained.

GC-MS/EI: 212

15 **Example 11**

Coupling of p-bromoacetophenone with phenol (catalyst from example 2)

- 1.35 g (6.7 mmol) of p-bromoacetophenone, 630 mg (6.7 mmol) of phenol,
20 1.8 g (13.5 mmol) of potassium carbonate and 360 mg (0.7 mmol) of the
catalyst from example 2 are stirred in 50 ml of dioxane under an argon
atmosphere at 110°C for 12 h. The GC analysis of the crude product
indicates product formation with 18% conversion.

Exempl 12**Coupling of p-bromoacetophenone with 2-hydroxypyridine**
(Catalyst from example 1)

- 5 1.35 g (6.7 mmol) of p-bromoacetophenone, 650 mg (6.7 mmol) of 2-hydroxypyridine, 1.8 g (13.5 mmol) of potassium carbonate and 300 mg (0.7 mmol) of the catalyst from example 1 are stirred in 50 ml of dioxane under an argon atmosphere at 110°C for 12 h. The reaction solution is
- 10 subsequently admixed with 40 ml of aqueous ammonia and extracted with ethyl acetate, and the combined organic extracts are dried under reduced pressure. After workup by column chromatography (hexane), 740 mg (52%) of a product mixture of N-4-acetylphenylpyridinone and 4-acetylphenoxy-2-pyridine (ratio 6:1) are obtained.
- 15 GC-MS/EI: 213 (M)

Example 13**Coupling of p-bromoacetophenone with 2-hydroxypyridine**
(Catalyst from example 2)

- 20 1.35 g (6.7 mmol) of p-bromoacetophenone, 650 mg (6.7 mmol) of o-hydroxypyridine, 1.8 g (13.5 mmol) of potassium carbonate and 360 mg (0.7 mmol) of the catalyst from example 2 are stirred in 50 ml of dioxane
- 25 under an argon atmosphere at 110°C for 12 h. The GC analysis of the crude product indicates product formation with 42% conversion.

Example 14**Coupling of p-bromoacetophenone with methanol (catalyst from example 1)**

- 5 1.34 g (13.5 mmol) of p-bromoacetophenone, 2 ml (20 mmol) of ethyl acetate, 17.5 ml of a 30% sodium methoxide solution and 300 mg (0.7 mmol) of the catalyst from example 1 are refluxed under an argon atmosphere for 12 h. The reaction solution is subsequently hydrolyzed
- 10 cautiously and extracted with dichloromethane, and the combined organic extracts are dried under reduced pressure. In addition to the product (35%) in a high proportion, the GC shows fragments of the aldol by-product. After workup by column chromatography (hexane), 820 mg (41%) of product were obtained.
- 15 GC-MS/EI: 150 (M)

Example 15**Coupling of p-bromoacetophenone with methanol (catalyst from example 2)**

- 20 1.34 g (13.5 mmol) of p-bromoacetophenone, 2 ml (20 mmol) of ethyl acetate, 17.5 ml of a 30% sodium methoxide solution and 360 mg (0.7 mmol) of the catalyst from example 2 are refluxed under an argon
- 25 atmosphere for 12 h. The reaction solution is subsequently hydrolysed cautiously and extracted with dichloromethane, and the combined organic extracts are dried under reduced pressure. In addition to the product (28%) in a high proportion, the GC shows fragments of the aldol by-product.
- 30

Example 16**Coupling of 4-iodotrifluoromethylbenzene with phenylacetylene**
(Catalyst from example 1)

5

2.7 g (10 mmol) of 4-iodotrifluoromethylbenzene, 1.3 g (12.5 mmol) of phenylacetylene, 2.2 g (20 mmol) of potassium tert-butoxide and 300 mg (0.7 mmol) of the catalyst from example 1 are stirred in 100 ml of dioxane under an argon atmosphere at 110°C for 21 h. The reaction solution is
10 filtered and dried under reduced pressure. After workup by column chromatography (hexane), 1.88 g (75%) of product are obtained.
GC-MS/EI: 246 (M)

Example 17

15

Coupling of 4-iodotrifluoromethylbenzene with phenylacetylene
(Catalyst from example 2)

2.7 g (10 mmol) of 4-iodotrifluoromethylbenzene, 1.3 g (12.5 mmol) of
20 phenylacetylene, 2.2 g (20 mmol) of potassium tert-butoxide and 360 mg (0.7 mmol) of the catalyst from example 2 are stirred in 100 ml of dioxane under an argon atmosphere at 110°C for 21 h. The GC analysis of the crude product indicates product formation (cf. with GC-MS/EI GZN 283-4) with 62% conversion.

25

Example 18**Coupling of p-tolyl iodide with phenylacetylene (catalyst from example 1)**

- 5 2.2 g (10 mmol) of p-tolyl iodide, 1.3 g (12.5 mmol) of phenylacetylene,
2.2 g (20 mmol) of potassium tert-butoxide and 300 mg (0.7 mmol) of the
catalyst from example 1 are stirred in 100 ml of dioxane under an argon
atmosphere at 110°C for 21 h. The reaction solution is filtered and dried
under reduced pressure. After workup by column chromatography
10 (hexane), 1.75 g (91%) of product are obtained.
GC-MS/EI: 192 (M)

Example 19

- 15 **Coupling of p-tolyl iodide with phenylacetylene (Catalyst from example 2)**

- 2.2 g (10 mmol) of p-tolyl iodide, 1.3 g (12.5 mmol) of phenylacetylene,
2.2 g (20 mmol) of potassium tert-butoxide and 360 mg (0.7 mmol) of the
catalyst from example 2 are stirred in 100 ml of dioxane under an argon
20 atmosphere at 110°C for 21 h. The GC analysis of the crude product
indicates product formation (cf. with GC-MS/EI GZN 277-8) with 50%
conversion.

Example 20

25

Coupling of bromobenzene with aniline (Catalyst from example 1)

- 1.05 g (6.7 mmol) of bromobenzene, 1.4 g (10.1 mmol) of caesium
carbonate and 300 mg (0.7 mmol) of the catalyst from example 1 are
30 stirred in 2 ml of aniline under an argon atmosphere at 170°C for 12 h.

The GC analysis of the crude product indicates the selective formation of the monoarylated compound GC-MS/EI: 169; (M)) and a conversion of 5%; triarylamine is not detected.

5 **Example 21**

Coupling of iodobenzene with aniline (Catalyst from example 1)

1.37 g (6.7 mmol) of iodobenzene, 1.4 g (10.1 mmol) of caesium
10 carbonate and 300 mg (0.7 mmol) of the catalyst from example 1 are
stirred in 2 ml of aniline under an argon atmosphere at 170°C for 12 h.
The GC analysis of the crude product indicates the selective formation of
the monoarylated compound (cf. with GC-MS/EI of 1126-7; triarylamine is
not detected. After workup by column chromatography, 810 mg (72%) of
15 product are obtained.

Example 22

Coupling of p-chloronitrobenzene with aniline (Catalyst from example 1)

20 1.06 g (6.7 mmol) of p-chloronitrobenzene, 1.4 g (10.1 mmol) of caesium
carbonate and 300 mg (0.7 mmol) of the catalyst from example 1 are
stirred in 2 ml of aniline under an argon atmosphere at 170°C for 12 h.
The GC analysis of the crude product indicates the selective formation of
25 the monoarylated compound and 83% conversion; triarylamine is not
detected. After workup by column chromatography, 1.14 g (80%) of
product are obtained.
GC-MS/EI: 214 (M)

Example 23**Coupling of iodobenzene with aniline (Catalyst from example 2)**

- 5 1.37 g (6.7 mmol) of iodobenzene, 1.4 g (10.1 mmol) of caesium carbonate and 360 mg (0.7 mmol) of the catalyst from example 2 are stirred in 2 ml of aniline under an argon atmosphere at 170°C for 12 h. The GC analysis of the crude product indicates the selective formation of the monoarylated compound and complete conversion; triarylamine is not
10 detected.

Example 24**Coupling of p-chloronitrobenzene with aniline (Catalyst from example 2)**

- 15 1.06 g (6.7 mmol) of p-chloronitrobenzene, 1.4 g (10.1 mmol) of caesium carbonate and 360 mg (0.7 mmol) of the catalyst from example 2 are stirred in 2 ml of aniline under an argon atmosphere at 170°C for 12 h. The GC analysis of the crude product indicates the selective formation of
20 the monoarylated compound (comparison with GC-MS/EI of 1101-7) and 91% conversion; triarylamine is not detected.

- Although the invention has been described in detail in the foregoing for the purpose of illustration, it is to be understood that such detail is solely for that
25 purpose and that variations can be made therein by those skilled in the art without departing from the spirit and scope of the invention except as it may be limited by the claims.